arXiv:math/0011067v1 [math.NT] 10 Nov 2000

2-EXTENSIONS WITH MANY POINTS

STÉPHAN SÉMIRAT

Abstract. We give defining equations for function fields over finite fields with many rational places. They are obtained from composita of quadratic extensions of the rational function field.

1. Introduction

We are interested in the following problem: given a prime power $q = p^e$ and a positive integer g, what is the maximal number of rational places $N_q(g)$ of a function field of genus g and finite constant field $\mathbb{F}_q$? This problem arise for instance in the theory of error correcting codes.

From the Riemann Hypothesis, we have the famous Hasse-Weil bound

$$N_q(g) \leq q + 1 + 2g\sqrt{q},$$

and J.P. Serre proved

$$N_q(g) \leq q + 1 + g[2\sqrt{q}],$$

where $[x]$ stands for the integer part of a real number x.

In fact, the Hasse-Weil bound is a particular case of the so-called "*méthode des formules explicites*", which majorates g for a fixed number of rational places. However, this method doesn't give the exact value of $N_q(g)$ and to this day, there is no method which could do so. We are thus lead to trying to reach or coming as close as possible to the best known upper bounds for $N_q(g)$, by way of finding (explicitely or not) function fields with the required parameters. The tables of G. van der Geer and M. van der Vlugt [8] record results on best known function fields for $1 \leq g \leq 50$ and q a small power of 2 or 3.

A classical method to obtain function fields with many rational places is given by class field theory. In this method, one has to manage with three parameters: a function field $K/\mathbb{F}_q$, a set S of rational places of K, and an effective divisor $\mathfrak{m}$ with support disjoint from S. Then the ray class field $K_S^{\mathfrak{m}}$ associated with $(K, S, \mathfrak{m})$ has its degree over K controlled by S (if S is sufficently large, one has $K_S^{m} = K$). The ramification of $K_S^{\mathfrak{m}}/K$ (and thus the genus) is controlled by $\mathfrak{m}$, and the number of rational places of $K_S^{\mathfrak{m}}$ is greater than $[K_S^{\mathfrak{m}} : K] \times |S|$, since all the places of S split totally in $K_S^{\mathfrak{m}}$. In particular, for S large and $\deg \mathfrak{m}$ small one can hope to obtain some "good" function fields. For instance, when $\mathfrak{m} = 0$, then $K_S^{\mathfrak{m}}$ is the Hilbert class field associated with (K, S) and if S equals the set of rational places of K, then the Hurwitz genus Formula gives

$$\frac{N(K_S^0)}{g_{K_S^0} - 1} \geq \frac{N(K)}{g_K - 1},$$

where we denote by $N(K)$ and g_K (resp. $N(K_S^0)$ and $g_{K_S^0}$ the number of rational places and genus of K (resp. K_S^0).

One may verify in the tables [8] that in general, when q is fixed, the quotient $N_q(g)/g$ decreases.

That is the greater $N(K)$ is, the smaller $h_S(K)$ should be, when S is chosen to be the whole set of rational places of K.
However, two problems arise with this method. The first one is that we don't know how to calculate the class number of a general pair (K, S). The second one is that for the applications, one needs explicit equations for the function fields involved. Such equations can hardly be found using general class field theory.

2. The method

In the following, we will apply the above class field method in it simplest form. We will construct some $(\mathbb{Z}/2\mathbb{Z})^n$-sub-fields of some ray class field associated with quadratic extensions of the rational function field $\mathbb{F}_q(x)$. The good thing with this is that we are able to give explicit equations for such fields.

Let us start with a simple example.

Example 2.1. *Let $k = \mathbb{F}_3(x)$ and let $K = k(y)$ be the quadratic extension of k defined by*

$$y^2 = f(x) = 2(x^3 + 2x + 2)(x^3 + 2x + 1).$$

The genus of $K/\mathbb{F}_3$ is $g_K = 2$. Now consider the compositum $L = K_1.K_2$ where $K_1 = k(y_1)$ and $K_2 = k(y_2)$ are defined by the equations

$$y_1^2 = f_1(x) = 2(x^3 + 2x + 2) \text{ and } y_2^2 = f_2(x) = x^3 + 2x + 1.$$

Since $f_1 f_2 = f$, K is a subfield of L and the ramification in L/K occurs only at the place of degree 2 above $(\frac{1}{x})$. The Hurwitz Formula gives then $g_L = 4$, for g_L the genus of L. Furthermore, since $f_i(0)$, $f_i(1)$, and $f_i(2)$ are all squares of $\mathbb{F}_3$, we have that $N(L) = 4 \times 3 = 12$, which is the maximal number of rational places for a function fields of genus 4 over $\mathbb{F}_3$.

Now we give more general facts.
With the help of the genus formulae for quadratic extensions (see for instance [6, Propositions VI.2.3 and VI.4.1], or [4]), the genus computation for a ramified 2-elementary extension of the rational function field can be done using the following theorem, which is a particular case of a theorem of G. Frey and H. G. Rück. A proof can be found in [2].

Theorem 2.2. *Let L/k be a Galois extension such that $\mathrm{Gal}(L/k) \simeq (\mathbb{Z}/2\mathbb{Z})^n$ and let K_I, $I \subset \{1, \ldots, n\}$ be the $2^n - 1$ sub-fields of L, quadratic over k, and let g_{K_I} be their respective genus. Then the genus g_L of L is given by*

$$g_L = \sum_i g(K_I)$$

Remarks 2.3.

- *In odd characteristic, we easily have*

$$g_L = 2^{n-1}\left(\sum_{P,\ e_P>1} \deg P - 4\right) + 1.$$

- *An analog formula is proven in* [7] *in the hard case of Artin-Schreier extension.*

For the applications, for insance to the geometric Goppa codes, one needs basis for some Riemann-Roch vector spaces. Such basis can be obtained as soon as one possesses a defining equation for the extension L/k (for instance, the software Kash [3] knows how to do it). For our extensions L/k, the following proposition gives us such an equation. Therefore, we are in the best situation for the applications.

Proposition 2.4. *Let L/k be the compositum of n disjoint quadratic extensions K_i/k given by $K_i = \mathbb{F}_q(x, y_i)$ with, for each i, y_i satisfying:*

- *in odd characteristic, $y_i^2 = f_i(x)$, where $f_i(x) \in \mathbb{F}_q[x]$ is square-free,*
- *in even characteristic, $y_i^2 + y_i = f_i(x)$, where $f_i(x) \in \mathbb{F}_q(x)$ such that for all $z \in \mathbb{F}_q(x)$, $f_i(x) \neq z^2 + z$.*

Then

$$y = \sum_i y_i \text{ (resp. } y = \prod_i y_i\text{)}$$

is a primitive element for L/k in odd (resp. even) characteristic.

Proof. The Galois group of L/k is generated by the σ_i, $1 \leq i \leq n$, such that in odd characteristic (resp. even) $\sigma_i(y_i) = -y_i$ (resp. $\sigma_i(y_i) = y_i + 1$), and, for $i \neq j$, $\sigma_i(y_j) = y_j$. The element

$$y = \sum_{i=1}^{n} y_i \text{ (resp. } y = \prod_{i=1}^{n} y_i\text{)},$$

is fixed by no element of $\mathrm{Gal}(L/k)$. Thus, $L = k(y)$.

□

Corollary 2.5. *Let $L = k(y_1, \ldots, y_n) = k(y)$, with y defined as in the above proposition. Then a minimal polynomial for y over k is given by*

$$P(Y, x) = \prod_I (Y - \sigma_I(y)) \in k[Y],$$

where for each $I \subseteq \{1, \ldots, n\}$,

$$\sigma_I(y) = \prod_{i=1..n,\ i \notin I} y_i \prod_{i=1..n,\ i \in I} \begin{cases} -y_i & \text{in odd characteristic,} \\ y_i + 1 & \text{in even characteristic.} \end{cases}$$

Example 2.6.

- *In odd characteristic,*
  - *the field $L = \mathbb{F}_q(x, y_1, y_2)$ with $y_i^2 = f_i(x)$ is given by $L = \mathbb{F}_q(x, y)$ with y satisfying*
    $$y^4 - 2(f_1 + f_2)y^2 + (f_1 - f_2)^2 = 0,$$
  - *the field $L = \mathbb{F}_q(x, y_1, y_2, y_3)$ with $y_i^2 = f_i(x)$ is given by $L = \mathbb{F}_q(x, y)$ with y satisfying*[1]

$$y^8 - 4\left(\sum_i f_i\right) y^6 + 2\left(3\sum_i f_i^2 + 2\sum_{i\neq j} f_f f_j\right) y^4 - 4\left(\sum_i f_i^3 - \sum_{i\neq j} f_i^2 f_j + 10 f_1 f_2 f_3\right) y^2$$
$$+ \sum_i f_i^4 - 4\sum_{i\neq j} f_i^3 f_j + 6\sum_{i\neq j} f_i^2 f_j^2 + 4\sum_{\substack{i\neq j,\ i\neq l \\ j\neq l}} f_i^2 f_j f_l = 0.$$

- *In even characteristic,*
  - *the field $L = \mathbb{F}_q(x, y_1, y_2)$ with $y_i^2 + y_i^2 = f_i(x)$ is given by $L = \mathbb{F}_q(x, y)$ with y satisfying*
    $$y^4 + y^3 + (f_1 + f_2)y^2 + f_1 f_2 y + f_1^2 f_2^2 = 0,$$

---

[1] The subscripts in the sums aren't ordered, for instance with $1 \leq i \leq 3$, the sum $\sum_{i\neq j} f_i f_j$ possesses 3 terms.

– *the field* $L = \mathbb{F}_q(x, y_1, y_2, y_3)$ *with* $y_i^2 + y_i = f_i(x)$ *is given* $L = \mathbb{F}_q(x, y)$ *with* y *satisfying*[1]

$$y^8 + y^7 + \left(\sum_i f_i\right) y^6 + \left(f_1f_2f_3 + \sum_{i \neq j} f_if_j\right) y^5 + \left(f_1f_2f_3 + \sum_{i\neq j} f_i^2 f_j^2\right) y^4$$

$$+ \left(f_1^2f_2^2f_3^2 + \sum_{\substack{i\neq j,\ i\neq l \\ j\neq l}} f_if_j^2f_l^2\right) y^3 + \left(\sum_{\substack{i\neq j,\ i\neq l \\ j\neq l}} f_i^3f_j^2f_l^2\right) y^2 + f_1^3f_2^3f_3^3 y + f_1^4f_2^4f_3^4.$$

3. TABLES

In the following tables, w denotes a generator for $\mathbb{F}_q^*$. We give in the last column either the exact value for $N_q(g)$, or a range $[a, b]$ (written $a - b$, following the notations in [8]) such that $N_q(g) \in [a, b]$ and there exists a function field defined over $\mathbb{F}_q$, with genus g and with at least a rational places.

We give the defining equations for function fields $L/\mathbb{F}_q$ with genus g_L obtained using the above method, and whose number of rational places $N(L)$ is near to, or reach, or is better than a. In the last case $N(L) > a$, estabilhing a new minoration for the corresponding value of $N_q(g)$, we use bold face.

The calculation have been made with the help of the algebraic number theory software Kash, v2.2 [3].

$\mathbf{q}$	$\mathbf{g_L}$	$\mathbf{f_i}$	$\mathbf{L = \mathbb{F}_q(x, y)}$	$\mathbf{N(L)}$	
2	1	$f_1 = \frac{1}{x}$ $f_2 = \frac{1}{x+1}$	$y^4 + y^3 + \frac{1}{x(x+1)}y^2 + \frac{1}{x(x+1)}y + \frac{1}{x^2(x+1)^2}$	4	5
	2	$f_1 = x$ $f_2 = x^3$	$y^4 + y^3 + (x^3 + x)y^2 + x^4y + x^8$	5	6
	3	$f_1 = \frac{1}{x}$ $f_2 = \frac{x}{x^2+x+1}$	$y^4 + y^3 + \frac{(x+1)^2}{x(x^2+x+1)}y^2 + \frac{1}{x(x^2+x+1)}y + \frac{1}{(x^2+x+1)^2}$	6	7
	4	$f_1 = x^3 + x$ $f^2 = x + 1/x$	$y^4 + y^3 + (x^4 + 1)/xy^2 + (x^4 + 1)y + x^8 + 1$	7	8
	5	$f_1 = x^3 + x$ $f_2 = x^5 + x$	$y^4+y^3+(x^5+x^3)y^2+(x^8+x^6+x^4+x^2)y+x^{16}+x^{12}+x^8+x^4$	9	9
	7	$f_1 = x^3 + x$ $f_2 = \frac{x(x+1)}{x^3+x+1}$	$y^4+y^3+\frac{x^3(x+1)(x^2+x+1)}{x^3+x+1}y^2+\frac{x^2(x+1)^3}{x^3+x+1}y+\frac{x^4(x+1)^6}{(x^3+x+1)^2}$	10	10
	9	$f_1 = \frac{x(x+1)}{x^3+x+1}$ $f_2 = \frac{x(x+1)}{x^3+x^2+1}$	$y^4 + y^3 + \frac{x^2(x+1)^2}{(x^3+x+1)(x^3+x^2+1)}y^2 + \frac{x^2(x+1)^2}{(x^3+x+1)(x^3+x^2+1)}y + \frac{x^4(x+1)^4}{(x^3+x+1)^2(x^3+x^2+1)^2}$	12	12
	11	$f_1 = \frac{x(x+1)}{x^3+x+1}$ $f_2 = \frac{x(x+1)}{x^4+x+1}$	$y^4 + y^3 + \frac{x^2(x+1)^2}{(x^3+x+1)(x^4+x+1)}y^2 + \frac{x^2(x+1)^2}{(x^3+x+1)(x^4+x+1)}y + \frac{x^4(x+1)^4}{(x^3+x+1)^2(x^4+x+1)^2}$	12	14
	15	$f_1 = x + \frac{x}{x^2+x+1}$ $f_2 = x+1+\frac{x+1}{x^2+x+1}$ $f_3 = x^3 + x$	$y^8 + y^7 + \frac{x^4(x+1)}{x^2+x+1}y^6 + \frac{x^2(x+1)^7}{(x^2+x+1)^2}y^5 + \frac{x^5(x+1)^5(x^4+x+1)}{(x^2+x+1)^4}y^4 + \frac{x^6(x+1)^{12}}{(x^2+x+1)^4}y^3 + \frac{x^{12}(x+1)^{11}}{(x^2+x+1)^5}y^2 + \frac{x^{12}(x+1)^{15}}{(x^2+x+1)^6}y + \frac{x^{16}(x+1)^{20}}{(x^2+x+1)^8}$	16	17

$\mathbf{q}$	$\mathbf{g_L}$	$\mathbf{f_i}$	$\mathbf{L = \mathbb{F}_q(x, y)}$	$\mathbf{N(L)}$	
	33	$f_1 = \frac{x(x+1)}{x^3+x+1}$ $f_2 = \frac{x(x+1)}{x^3+x^2+1}$ $f_3 = \frac{x(x+1)}{x^4+x+1}$	$y^8 + y^7 + \frac{x(x+1)(x^2+x+1)}{(x^3+x+1)(x^3+x^2+1)(x^4+x+1)}y^6 + \frac{x^2(x+1)^2}{(x^3+x+1)(x^3+x^2+1)}y^5 + \frac{x^3(x+1)^3(x^4+x^3+1)(x^4+x^3+x^2+x+1)}{(x^3+x+1)(x^3+x^2+1)(x^4+x+1)}y^4 + (x^3+x+1)^2(x^3+x^2+1)^2(x^4+x+1)y^3 + \frac{x^7(x+1)^7(x^2+x+1)^2}{(x^3+x+1)^3(x^3+x^2+1)^3(x^4+x+1)^3}y^2 + \frac{x^9(x+1)^9}{(x^3+x+1)^3(x^3+x^2+1)^3(x^4+x+1)^3}y + \frac{x^{12}(x+1)^{12}}{(x^3+x+1)^4(x^3+x^2+1)^4(x^4+x+1)^4}$	24	$28-29$
	41	$f_1 = x + \frac{x}{x^2+x+1}$ $f_2 = x+1+\frac{x+1}{x^2+x+1}$ $f_3 = x^3 + x$ $f_4 = \frac{x(x+1)}{x^3+x+1}$		32	$33-35$
4	1	$f_1 = x,\ f_2 = \frac{1}{x}$	$y^4 + y^3 + \frac{(x+1)^2}{x}y^2 + y + 1$	8	9
	2	$f_1 = x,\ f_2 = x^3$	$y^4 + y^3 + (x^3+x)y^2 + x^4y + x^8$	9	10
	3	$f_1 = \frac{1}{x} + \frac{w}{x+w^2}$ $f_2 = \frac{1}{x} + \frac{w^2}{x+w}$	$y^4 + y^3 + \frac{x+1}{(x+w)(x+w^2)}y^2 + \frac{(x+1)^2}{x^2(x+w)(x+w^2)}y + \frac{(x+1)^4}{x^4(x+w)^2(x+w^2)^2}$	14	14
	4	$f_1 = x^3+1\ f_2 = x+1/x$	$y^4+y^3+\frac{(x+1)(x^3+x^2+1)}{x}y^2+\frac{(x+1)^3(x+w)(x+w^2)}{x}y+(x+1)^6(x+w)^2(x+w^2)^2$	15	15
	5	$f_1 = x^3+1,\ f_2 = x^5+x$	$y^4+y^3+(x+1)(x^2+wx+w)(x^2+w^2x+w^2)y^2+x(x+1)^5(x+w)(x+w^2)y+x^2(x+1)^{10}(x+w)^2(x+w^2)^2$	17	$17-18$
	7	$f_1 = x^3+1$ $f_2 = \frac{x(x+w^2)}{x^3+w^2}$	$y^4 + y^3 + (x^6+wx^3+x^2+w^2x+w^2)/(x^3+w^2)y^2 + (x^5+w^2x^4+x^2+w^2x)/(x^3+w^2)y+(x^{10}+wx^8+x^4+wx^2)/(x^6+w)$	18	$21-22$

$\mathbf{q}$	$\mathbf{g_L}$	$\mathbf{f_i}$	$\mathbf{L} = \mathbb{F}_{\mathbf{q}}(\mathbf{x}, \mathbf{y})$	$\mathbf{N(L)}$	
	11	$f_1 = x^3 + 1$ $f_2 = x + \frac{1}{x}$ $f_3 = x + 1 + \frac{1}{x+1}$	$y^8 + y^7 + \frac{(x+w)(x+w^2)(x^3+x+1)}{x(x+1)}y^6 + \frac{(x^3+wx^2+1)(x^3+w^2x^2+1)}{x}y^5 + \frac{(x^3+x^2+1)(x^5+x^2+1)}{x^2}y^4 + (x+1)^2(x+w)^2(x+w^2)(x^3+wx^2+1)(x^3+w^2x^2+1)y^3 + x(x+1)^3(x+w)^3(x+w^2)^3(x^3+x+1)y^2 + xx^3(x+1)^6(x+w)^3(x+w^2)^3y + x^4(x+1)^8(x+w)^4(x+w^2)^4$	26	$26 - 30$
8	1	$f_1 = \frac{1}{x}, \ f_2 = \frac{w}{x+w^4}$	$y^4 + y^3 + (w^3x + w^4)/(x^2 + w^4x)y^2 + w/(x^2 + w^4x)y + w^2/(x^4 + wx^2)$	14	14
	2	$f_1 = \frac{1}{x} + \frac{w^6}{x+1}$ $f_2 = \frac{w^3}{x}$	$y^4 + y^3 + \frac{w^5(x+w^3)}{x(x+1)}y^2 + \frac{w^5(x+w^5)}{x^2(x+1)}y + \frac{w^3(x+w^5)^2}{x^4(x+1)^2}$	17	18
	3	$f_1 = \frac{1}{x} + \frac{w}{x+w^4}, \ f_2 = \frac{1}{x} + \frac{w^5}{x+w^6}$	$y^4 + y^3 + \frac{w^6(x+1)}{(x+w^4)(x+w^6)}y^2 + \frac{(x+w)(x+w^2)}{x^2(x+w^4)(x+w^6)}y + \frac{(x+w)^2(x+w^2)^2}{x^4(x+w^4)(x+w^6)}$	24	24
	4	$f_1 = x^3 + x + 1$ $f_2 = x + \frac{1}{x+1}$	$y^4 + y^3 + \frac{x(x+w^3)(x+w^5)(x+w^6)}{x+1}y^2 + \frac{(x+w)(x+w^2)(x+w^4)(x^2+x+1)}{x+1}y + \frac{(x+w)^2(x+w^2)^2(x+w^4)^2(x^2+x+1)^2}{(x+1)^2}$	25	$25 - 27$
	5	$f_1 = \frac{1}{x^2+x+1}$ $f_2 = \frac{w(x+w^6)}{x^2+w^6x+w}$	$y^4 + y^3 + (wx^3 + wx^2 + w^4x + w^3)/(x^4 + w^2x^3 + w^4x^2 + w^5x + w)y^2 + (wx+1)/(x^4 + w^2x^3 + w^4x^2 + w^5x + w)y + (w^2x^2 + 1)/(x^8 + w^4x^6 + wx^4 + w^3x^2 + w^2)$	28	$29 - 32$
	7	$f_1 = \frac{1}{x} + \frac{w(x+w^3)}{x^2+w^5x+w}$ $f_2 = \frac{1}{x} + \frac{w^2(x+w^6)}{x^2+w^3x+w^2}$	$y^4 + y^3 + (w^4x^3 + w^3x^2 + w^2x + 1)/(x^4 + w^2x^3 + w^2x^2 + w^5x + w^3)y^2 + (w^2x^4 + w^4x^3 + w^5x^2 + w^4x + w^3)/(x^6 + w^2x^5 + w^2x^4 + w^5x^3 + w^3x^2)y + (w^4x^8 + wx^6 + w^3x^4 + wx^2 + w^6)/(x^{12} + w^4x^{10} + w^4x^8 + w^3x^6 + w^6x^4)$	**34**	$33 - 39$

$\mathbf{q}$	$\mathbf{g_L}$	$\mathbf{f_i}$	$\mathbf{L} = \mathbb{F}_{\mathbf{q}}(\mathbf{x},\mathbf{y})$	$\mathbf{N(L)}$	
	8	$f_1 = 1/x + 1/(x+w^2)$ $f_2 = w/(x+w^2) + w^6/(x+w^6)$ $f_3 = w^5/(x+w^2)$	$y^8 + y^7 + (x+w)/(x^3+x^2+wx)y^6 + (w^3x^2+w^6x+w^4)/(x^4+x^3+w^2x^2+w^4x+w^5)y^5 + (wx^3+w^5x^2+w^3x+1)/(x^7+w^2x^6+x^5+w^2x^4+w^2x^3+w^4x^2)y^4 + (wx^3+w^3x^2+1)/(x^9+x^7+w^4x^5+wx^3+w^3x)y^3 + (w^3x^3+w^4x^2+w^6x+1)/(x^{13}+x^{12}+w^3x^{11}+x^{10}+w^2x^9+w^4x^8+w^6x^7+wx^6+w^5x^5+w^3x^4+w^4x^3)y^2 + (wx^3+w^6x^2+w^4x+w^2)/(x^{15}+x^{14}+w^6x^{13}+w^5x^{12}+w^6x^{11}+w^2x^7+w^2x^6+wx^5+x^4+wx^3) * y + (w^6x^4+w^5)/(x^{20}+x^{16}+wx^{12}+w^2x^8+w^6x^4)$	33	$34-43$
	11	$f_1 = x^3+x+1$ $f_2 = x + \dfrac{1}{x+1}$ $f_3 = x+1+\dfrac{1}{x}$	$y^8 + y^7 + (x^5+x^4+x^3+x^2+1)/(x^2+x)y^6 + (x^7+x^5+x^4+x+1)/(x^2+x)y^5 + (x^8+x^6+x^5+x^3+x^2+x+1)/(x^2+x)y^4 + (x^{14}+x^{12}+x^8+x^7+x^5+x^3+1)/(x^4+x^2)y^3 + (x^{19}+x^{18}+x^{17}+x^{16}+x^{14}+x^{13}+x^{12}+x^{11}+x^{10}+x^9+x^4+x^3+1)/(x^6+x^5+x^4+x^3)y^2 + (x^{21}+x^{18}+x^{16}+x^{14}+x^{12}+x^{11}+x^{10}+x^7+x^5+x^4+x^3+x+1)/(x^6+x^5+x^4+x^3)y + (x^{28}+x^{16}+x^8+x^4+1)/(x^8+x^4)$	48	$48-54$
16	1	$f_1 = \dfrac{1}{x},\ f_2 = \dfrac{1}{x+w^5}$	$y^4+y^3+w^5/(x^2+w^5x)y^2+1/(x^2+w^5x)y+1/(x^4+w^{10}x^2)$	24	25
	2	$f_1 = x^3+x+1$ $f_2 = w^5x$	$y^4+y^3+(x^3+w^{10}x+1)y^2+(w^5x^4+w^5x^2+w^5x)y+w^{10}x^8+w^{10}x^4+w^{10}x^2$	33	33
	3	$f_1 = \dfrac{1}{x}+\dfrac{1}{x+w^{10}}$ $f_2 = \dfrac{1}{x}+\dfrac{w^5}{x+w^5}$	$y^4+y^3+(w^{10}x+w^{10})/(x^2+x+1)y^2+w^5/(x^3+w^5x^2)y+w^{10}/(x^6+w^{10}x^4)$	38	38
	4	$f_1 := \frac{w^7}{x}+\frac{w}{x+w^{14}}$ $f_2 = \frac{w^2}{x}+\frac{1}{x+w^6}$	$y^4+y^3+(w^6x^2+w^2x+w^2)/(x^3+w^8x^2+w^5x)y^2+(w^7x^2+wx+w^{14})/(x^4+w^8x^3+w^5x^2)y+(w^{14}x^4+w^2x^2+w^{13})/(x^8+wx^6+w^{10}x^4)$	45	$45-46$
	5	$f_1 = x^3+x$ $f_2 = x^5$	$y^4+y^3+(x^5+x^3+x)y^2+(x^8+x^6)y+x^{16}+x^{12}$	49	$49-54$
32	1	$f_1 = \dfrac{1}{x},\ f_2 = \dfrac{1}{x+1}$	$y^4+y^3+y^2/(x^2+x)+y/(x^2+x)+1/(x^4+x^2)$	44	44
	3	$f_1 = \dfrac{1}{x}+\dfrac{1}{x+1}$ $f_2 = \dfrac{1}{x}+\dfrac{w}{x+w^9}$	$y^4 + y^3 + \dfrac{w^{18}(x+w^3)}{(x+1)(x+w^9)}y^2 + \dfrac{w^{18}(x+w^{22})}{x^2(x+1)(x+w^9)} + \dfrac{w^5(x+w^{22})}{x^4(x+1)^2(x+w^9)^2}$	**64**	$63-64$
	5	$f_1 = \dfrac{1}{x^2+w^3x+w}$ $f_2 = \dfrac{x+w^{24}}{x^2+w^{27}x+w^{10}}$	$y^4+y^3+(x^3+w^{26}x^2+wx+w^3)/(x^4+w^{18}x^3+x^2+w^6x+w^{11})y^2+(x+w^{24})/(x^4+w^{18}x^3+x^2+w^6x+w^{11})y+(x^2+w^{17})/(x^8+w^5x^6+x^4+w^{12}x^2+w^{22})$	76	$83-86$

q	$\mathbf{g_L}$	$\mathbf{f_i}$	$\mathbf{L} = \mathbb{F}_\mathbf{q}(\mathbf{x},\mathbf{y})$	$\mathbf{N(L)}$	
	7	$f_1 = \frac{1}{x} + \frac{1}{x+1}$ $f_2 = \frac{1}{x} + \frac{w^{11}}{x+w^{21}}$ $f_3 = \frac{w^{21}}{x}$	$y^8 + y^7 + (w^{24}x^2 + w^{11})/(x^3 + w^{25}x^2 + w^{21}x)y^6 + (w^9x^3 + w^{18}x^2 + wx + w^{11})/(x^5 + w^{25}x^4 + w^{21}x^3) * y^5 + (w^{25}x^4 + w^{23}x^3 + w^{30}x^2 + wx + w^{11})/(x^8 + w^{19}x^6 + w^{11}x^4)y^4 + (w^{18}x^4 + w^{11}x^3 + w^{21}x^2 + wx + w^{22})/(x^{10} + w^{19}x^8 + w^{11}x^6)y^3 + (w^{11}x^4 + w^{28}x^2 + w^2)/(x^{13} + w^{25}x^{12} + w^{24}x^{11} + w^{13}x^{10} + w^{14}x^9 + w^5x^8 + wx^7)y^2 + (w^{27}x^3 + w^{29}x^2 + x + w^2)/(x^{15} + w^{25}x^{14} + w^{24}x^{13} + w^{13}x^{12} + w^{14}x^{11} + w^5x^{10} + wx^9)y + (w^5x^4 + w^{13})/(x^{20} + w^7x^{16} + w^{22}x^{12})$	**98**	$90 - 108$
	9	$f_1 = \frac{1}{x} + \frac{1}{x+1}$ $f_2 = \frac{1}{x} + \frac{w}{x+w^9}$ $f_3 = \frac{w^{24}}{x+w^5}$	$y^8 + y^7 + (w^{14}x^2 + w^8x + w^{17})/(x^3 + w^{24}x^2 + wx + w^{14})y^6 + (w^{11}x^3 + w^{24}x^2 + w^{30}x + w^{25})/(x^5 + w^{24}x^4 + wx^3 + w^{14}x^2)y^5 + (w^{30}x^6 + w^7x^5 + w^6x^4 + w^{10}x^3 + w^{11}x^2 + w^{28})/(x^{10} + w^{17}x^8 + w^2x^6 + w^{28}x^4)y^4 + (w^{22}x^4 + w^{20}x^3 + w^{19}x^2 + w^{11}x + w^{27})/(x^{10} + w^{17}x^8 + w^2x^6 + w^{28}x^4)y^3 + (w^5x^4 + w^{30}x^3 + w^{12}x^2 + w^{12}x + w^{21})/(x^{13} + w^{24}x^{12} + w^{10}x^{11} + w^{20}x^{10} + w^{11}x^9 + w^{28}x^8 + w^{24}x^7 + w^{18}x^6 + w^{29}x^5 + w^{11}x^4)y^2 + (w^2x^3 + w^{24}x^2 + w^15x + w^6)/(x^{15} + w^{24}x^{14} + w^{10}x^{13} + w^{20}x^{12} + w^{11}x^{11} + w^{28}x^{10} + w^{24}x^9 + w^{18}x^8 + w^{29}x^7 + w^{11}x^6)y + (w^{13}x^4 + w^8)/(x^{20} + w^3x^{16} + w^4x^{12} + w^{25}x^8)$	104	$108 - 130$
	11	$f_1 = \frac{1}{x} + \frac{1}{x+1}$ $f_2 = \frac{1}{x} + \frac{w^4}{x+w^2}$ $f_3 = \frac{w^7}{x} + \frac{w^{13}}{x+w^{22}}$	$y^8 + y^7 + (w^{25}x^3 + w^{30}x^2 + w^2x + 1)/(x^4 + w^4x^3 + w^{23}x^2 + w^{24}x)y^6 + (w^{13}x^4 + w^8x^3 + w^{15}x^2 + w^{28}x + 1)/(x^6 + w^4x^5 + w^{23}x^4 + w^{24}x^3)y^5 + (w^{27}x^6 + w^{16}x^5 + w^{27}x^4 + w^8x^3 + x^2 + w^{24}x + w^{17})/(x^{10} + w^8x^8 + w^{15}x^6 + w^{17}x^4)y^4 + (w^{26}x^6 + w^{18}x^5 + w^8x^4 + w^{15}x^3 + wx^2 + w^{24}x + 1)/(x^{12} + w^8x^{10} + w^{15}x^8 + w^{17}x^6)y^3 + (w^{20}x^7 + w^{25}x^6 + w^{26}x^5 + w^{30}x^4 + w^7x^3 + w^{21}x^2 + w^2x + 1)/(x^{16} + w^4x^{15} + wx^{14} + w^4x^{13} + w^{24}x^{12} + w^2x^{11} + w^{11}x^{10} + w^{22}x^9 + w^9x^8 + w^{10}x^7)y^2 + (w^8x^6 + w^{29}x^5 + w^{10}x^4 + w^9x^3 + w^{28}x^2 + w^3x + 1)/(x^{18} + w^4x^{17} + wx^{16} + w^4x^{15} + w^{24}x^{14} + w^2x^{13} + w^{11}x^{12} + w^{22}x^{11} + w^9x^{10} + w^{10}x^9)y + (w^{21}x^8 + w^{12}x^4 + 1)/(x^{24} + w^{16}x^{20} + w^{30}x^{16} + w^3x^{12})$	**120**	$113 - 152$
64	1	$f_1 = \frac{1}{x},\ f_2 = \frac{1}{x+w^{27}}$	$y^4 + y^3 + w^{27}/(x^2 + w^{27}x)y^2 + 1/(x^2 + w^{27}x)y + 1/(x^4 + w^{54}x^2)$	80	81
	3	$f_1 = \frac{1}{x^2 + w^{25}x + 1}$ $f_2 = \frac{w^{14}}{x+w^{28}}$	$y^4 + y^3 + (w^{14}x^2 + w^{17}x + w^{30})/(x^3 + w^{38}x^2 + w^{50}x + w^{28})y^2 + w^{14}/(x^3 + w^{38}x^2 + w^{50}x + w^{28})y + w^{28}/(x^6 + w^{13}x^4 + w^{37}x^2 + w^{56})$	104	113

$\mathbf{q}$	$\mathbf{g_L}$	$\mathbf{f_i}$	$\mathbf{L} = \mathbb{F}_\mathbf{q}(\mathbf{x}, \mathbf{y})$	$\mathbf{N(L)}$	
	5	$f_1 = \frac{1}{x^2 + w^{11}x + 1}$ $f_2 = \frac{1}{x^2 + w^{25}x + 1}$	$y^4 + y^3 + w^{27}x/(x^4 + w^{27}x^3 + w^{36}x^2 + w^{27}x + 1)y^2 + 1/(x^4 + w^{27}x^3 + w^{36}x^2 + w^{27}x + 1)y + 1/(x^8 + w^{54}x^6 + w^9x^4 + w^{54}x^2 + 1)$	128	$130 - 145$
128	1	$f_1 = \frac{1}{x}$, $f_2 = \frac{w^{11}}{x + w^{111}}$	$y^4 + y^3 + (w^{87}x + w^{111})/(x^2 + w^{111}x)y^2 + w^{11}/(x^2 + w^{111}x)y + w^{22}/(x^4 + w^{95}x^2)$	150	150
	3	$f_1 = \frac{1}{x} + \frac{w}{x + w^{19}}$ $f_2 = \frac{1}{x} + \frac{w^{74}}{x + w^{79}}$	$y^4 + y^3 + (w^{79}x + w^8)/(x^2 + w^{16}x + w^{98})y^2 + (w^{99}x^2 + w^{64}x + w^{98})/(x^4 + w^{16}x^3 + w^{98}x^2)y + (w^{71}x^4 + wx^2 + w^{69})/(x^8 + w^{32}x^6 + w^{69}x^4)$	**192**	$191 - 192$
	5	$f_1 = \frac{1}{x}$ $f_2 = \frac{w^{11}}{x + w^{111}}$ $f_3 = \frac{w^{42}}{x + w^{63}}$	$y^8 + y^7 + (w^{110}x^2 + w^{64}x + w^{47})/(x^3 + w^{55}x^2 + w^{47}x)y^6 + (w^{92}x + 1)/(x^3 + w^{55}x^2 + w^{47}x)y^5 + (w^{53}x^3 + w^{97}x^2 + w^{100}x + w^{36})/(x^6 + w^{110}x^4 + w^{94}x^2)y^4 + (w^{18}x + w^{53})/(x^6 + w^{110}x^4 + w^{94}x^2)y^3 + (w^{89}x^2 + w^{43}x + w^{26})/(x^9 + w^{55}x^8 + w^{50}x^7 + w^{38}x^6 + w^{97}x^5 + w^{22}x^4 + w^{14}x^3)y^2 + w^{32}/(x^9 + w^{55}x^8 + w^{50}x^7 + w^{38}x^6 + w^{97}x^5 + w^{22}x^4 + w^{14}x^3)y + w^{85}/(x^{12} + w^{93}x^8 + w^{61}x^4)$	216	$227 - 239$
	9	$f_1 = \frac{1}{x} + \frac{w}{x + w^{19}}$ $f_2 = \frac{1}{x} + \frac{w^{74}}{x + w^{79}}$ $f_3 = \frac{1}{x} + \frac{w^{80}}{x + w^{61}}$	$y^8 + y^7 + (w^{34}x^3 + w^{24}x^2 + w^{39}x + w^{32})/(x^4 + w^{84}x^3 + w^7x^2 + w^{32}x)y^6 + (w^{122}x^4 + w^{33}x^3 + w^{112}x^2 + w^{33}x + w^{32})/(x^6 + w^{84}x^5 + w^7x^4 + w^{32}x^3)y^5 + (w^{74}x^7 + w^{109}x^6 + w^{35}x^5 + w^{51}x^4 + w^{15}x^3 + w^{63}x^2 + w^{64}x + w^{64})/(x^{10} + w^{41}x^8 + w^{14}x^6 + w^{64}x^4)y^4 + (w^{69}x^7 + w^2x^6 + w^{30}x^5 + w^6x^4 + w^{47}x^3 + w^{116}x^2 + w^{64}x + w^{64})/(x^{12} + w^{41}x^{10} + w^{14}x^8 + w^{64}x^6)y^3 + (w^{55}x^9 + w^{45}x^8 + w^{118}x^7 + w^{35}x^6 + w^{66}x^5 + w^{95}x^4 + x^3 + w^8x^2 + w^{103}x + w^{96})/(x^{16} + w^{84}x^{15} + w^{93}x^{14} + w^{84}x^{13} + w^{100}x^{12} + w^{51}x^{11} + w^{38}x^{10} + w^{126}x^9 + w^{71}x^8 + w^{96}x^7)y^2 + (w^{95}x^9 + w^{53}x^8 + w^6x^7 + w^{70}x^6 + w^{98}x^5 + w^{98}x^4 + w^{47}x^3 + w^{98}x^2 + w^{103}x + w^{96})/(x^{18} + w^{84}x^{17} + w^{93}x^{16} + w^{84}x^{15} + w^{100}x^{14} + w^{51}x^{13} + w^{38}x^{12} + w^{126}x^{11} + w^{71}x^{10} + w^{96}x^9)y + (w^{42}x^{12} + wx^8 + w^{29}x^4 + w)/(x^{24} + w^{82}x^{20} + w^{28}x^{16} + wx^{12})$	**288**	$258 - 327$
3	1	$f_1 = x$ $f_2 = x^2 + x + 2$	$y^4 + (x^2 + 2x + 2)y^2 + x^4 + x^2 + 1$	6	7
	2	$f_1 = x$ $f_2 = x^3 + 2x + 1$	$y^4 + (x^3 + 1)y^2 + x^6 + 2x^4 + 2x^3 + x^2 + 2x + 1$	8	8
	3	$f_1 = (x + 1)(x^3 + x^2 + x + 2)$ $f_2 = x^2 + x + 2$	$y^4 + (x^4 + 2x^3 + x + 1)y^2 + x^8 + x^7 + 2x^5 + x^3 + x^2$	8	10

q	$\mathbf{g_L}$	$\mathbf{f_i}$	$\mathbf{L = \mathbb{F}_q(x, y)}$	$\mathbf{N(L)}$	
	4	$f_1 = x^3 + 2x + 1$ $f_2 = 2(x^3 + 2x + 2)$	$y^4 + 2y^2 + x^6 + x^4 + x^2$	12	12
	5	$f_1 = (x + 1)(x^3 + x^2 + 2x + 1)$ $f_2 = x^4 + x^3 + x^2 + 1$	$y^4 + (2x^4 + x^2 + 2)y^2 + x^6 + x^5 + x^4$	12	$12 - 13$
	6	$f_1 = x^3 + 2x + 1$ $f_2 = x^5 + 2x + 1$	$y^4 + (x^5 + x^3 + x + 2)y^2 + x^{10} + x^8 + x^6$	14	$14 - 15$
	7	$f_1 = (x^2 + 1)(x^2 + x + 2)(x^2 + 2x + 2)$ $f_2 = (x^2 + 1)(x^4 + x^3 + 2x + 1)$	$y^4 + (2x^6 + x^5 + 2x^4 + 2x^2 + 2x + 2)y^2 + x^{10} + x^6 + x^2$	16	16
	8	$f_1 = x^5 + 2x + 1$ $f_2 = x^5 + x^3 + x + 1$	$y^4 + (2x^5 + x^3 + 2)y^2 + x^6 + x^4 + x^2$	14	$15 - 18$
	9	$f_1 = x^2 + 1$ $f_2 = (x^2 + x + 2)(x^2 + 2x + 2)$ $f_3 = (x+1)(x+2)(x^2 + x + 2)$	$y^8 + (x^4 + 2x^3 + x^2 + x)y^6 + (x^8 + x^7 + 2x^4 + x^3 + x^2 + x) * y^4 + (2x^{10} + 2x^9 + x^8 + x^7 + 2x^6 + 2x^4 + 2x^3 + x + 2)y^2 + x^6 + x^4 + x^2$	16	19
	13	$f_1 = (x + 1)(x^3 + x^2 + 2x + 1)$ $f_2 = (x + 1)(x^2 + 1)$ $f_3 = x^3 + 2x + 1$	$y^8 + (2x^4 + 2x^3 + 2x^2)y^6 + (2x^7 + 2x^4 + x^3 + x^2)y^4 + (2x^{12} + 2x^{11} + x^{10} + x^9 + 2x^7 + x^5 + 2)y^2 + x^{16} + x^{13} + x^{12} + x^{11} + 2x^{10} + 2x^9 + x^7 + 2x^5 + x^4$	24	$24 - 25$
	17	$f_1 = (x + 2)(x^3 + x^2 + x + 2)$ $f_2 = x^4 + x^3 + x^2 + x + 1$ $f_3 = x^4 + 2x^3 + x^2 + 1$	$y^8 + (x^2 + x)y^6 + (x^5 + 2x^2 + x)y^4 + (2x^{12} + 2x^{10} + x^9 + 2x^6 + 2x^5 + x^3 + 2x^2 + x + 2)y^2 + x^{10} + x^9 + x^6 + 2x^3 + x^2$	24	$24 - 30$
	19	$f_1 = (x^2 + 1)(x^2 + x + 2)(x^2 + 2x + 2)$ $f_2 = (x^2 + 1)(x^3 + x^2 + 2x + 1)$ $f_3 = x^3 + 2x + 1$	$y^8 + (2x^6 + 2x^5 + x^4 + 2x^3 + 2x)y^6 + (x^{11} + x^{10} + 2x^9 + x^8 + x^7 + x^6 + 2x^5 + 2x^4 + x^3 + x^2 + 2x)y^4 + (2x^{18} + x^{17} + 2x^{16} + x^{15} + 2x^{14} + 2x^{12} + x^{11} + 2x^7 + x^6 + 2x^5 + 2x^4 + x^2 + 2x + 2)y^2 + x^{24} + 2x^{23} + x^{21} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + 2x^{15} + 2x^{14} + x^{13} + 2x^{10} + x^7 + 2x^6 + x^4 + x^2$	28	$28 - 32$
	21	$f_1 = (x^2 + 1)(x^2 + x + 2)(x^2 + 2x + 2)$ $f_2 = (x^2 + 1)(x^4 + x^3 + 2x + 1)$ $f_3 = (x^2 + x + 2)(x^4 + x + 2)$	$y^8 + (x^5 + 2x^4 + 2x^3 + x^2)y^6 + (x^{11} + x^9 + x^8 + 2x^7 + 2x^5 + 2x^4)y^4 + (2x^{18} + x^{17} + 2x^{15} + x^{12} + x^{11} + 2x^9 + x^6 + 2x^5 + x^3 + 2x^2 + 2)y^2 + x^{22} + x^{21} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{12} + x^{11} + 2x^9 + x^8$	32	$32 - 35$

$\mathbf{q}$	$\mathbf{g_L}$	$\mathbf{f_i}$	$\mathbf{L = \mathbb{F}_q(x, y)}$	$\mathbf{N(L)}$	
9	1	$f_1 = x(x + w^6)$ $f_2 = (x + w^2)$	$y^4 + (x^2 + w^5x + w^2)y^2 + x^4 + w^7x^3 + w^5x + 2$	16	16
	2	$f_1 = x$ $f_2 = (x+1)(x+w^5)(x+w^7)$	$y^4 + (x^3 + 2 * x + 2) * y^2 + x^6 + x^3 + 1$	18	20
	3	$f_1 = x(x + w)(x + 1)(x + w^3)$ $f_2 = x(x + w)(x + w^6)(x + w^7)$	$y^4 + (2x^4 + wx^3 + w^3x)y^2 + w^6x^6 + x^4 + w^2x^2$	28	28
	4	$f_1 = x^3 + w^2x^2 + w^2$ $f_2 = x^3 + w^6x^2 + w^6$	$y^4 + 2x^3y^2 + 2x^4 + x^2 + 2$	30	30
	5	$f_1 = x(x + w)$ $f_2 = (x + 1)(x + w^3)$ $f_3 = (x + w^6)(x + w^7)$	$y^8 + w^6y^6 + y^4 + (2x^6 + w^2x^4 + w^3x^3 + x^2 + w^5x + w^6)y^2 + 1$	32	$32 - 36$
	6	$f_1 = x(x^2 + w)(x^3 + wx^2 + 2x + w^7)$ $f_2 = x(x^2 + w)(x^3 + w^3x^2 + w^6x + w^3)$	$y^4 + (2x^6 + x^5 + w^2x^4 + wx^3 + 2x^2)y^2 + 2x^{10} + w^3x^9 + w^2x^8 + 2x^7 + x^6 + wx^5 + 2x^4 + w^2x^3 + x^2$	34	$35 - 40$
	7	$f_1 = (x^2 + w^2x + w^6)(x^4+wx^2+wx+w^2)$ $f_2 = (x^2 + w^2x + w^6)(x^4 + w^7x^2 + w^5x + w^6)$	$y^4 + (2x^6 + w^6x^5 + x^3 + 2x^2)y^2 + x^8 + 2x^7 + w^2x^6 + w^3x^5 + w^5x^4 + wx^3 + w^6x^2 + w^6x + 1$	36	$40 - 43$
	8	$f_1 = (x^3 + w^2x^2 + 2x + w^2)(x^5 + 2x + w^2)$ $f_2 = (x^3 + w^6x^2 + x + w^2)(x^5 + 2x + w^2)$	$y^4 + (2x^8 + w^6x^5 + x^4 + w^6x^3 + w^2x + 1)y^2 + 2x^{14} + w^2x^{13} + x^{12} + 2x^{10} + w^6x^9 + 2x^8 + w^6x^7 + 2x^6 + x^4 + 2x^2$	**40**	$38 - 47$
	9	$f_1 = w * x^4 + w^7,$ $f_2 = x^4 + x^3 + w^5 * x + w^6$ $f_3 = x^4 + w^3 * x^3 + w * x^2 + 2 * x$	$y^8 + (w^3x^4 + w^2x^3 + w^5x^2 + w^2x + 2)y^6 + (w^3x^8 + x^7 + w^7x^6 + w^2x^5 + w^6x^4 + w^6x^3 + w^5x^2 + x + w^5)y^4 + (w^2x^{12} + wx^{11} + w^6x^9 + w^2x^8 + w^3x^7 + w^2x^6 + wx^5 + w^2x^4 + w^7x^3 + w^2x^2 + w^7x + w^6)y^2 + x^{16} + w^3x^{15} + w^5x^{13} + w^6x^{12} + w^6x^{11} + wx^{10} + w^2x^9 + w^5x^8 + wx^7 + 2x^6 + wx^5 + w^3x^4 + w^6x^3 + w^6x^2 + w^7x + 2$	40	$48 - 51$

q	$\mathbf{g_L}$	$\mathbf{f_i}$	$\mathbf{L} = \mathbb{F}_\mathbf{q}(\mathbf{x}, \mathbf{y})$	$\mathbf{N}(\mathbf{L})$	
	17	$f_1 = (x^3 + wx^2 + wx + w^6)(x^3 + w^3x^2 + w^3x + w^2)$ $f_2 = (x^3 + wx^2 + wx + w^6)(x^3 + x + w^2)$ $f_3 = (x^3 + wx^2 + wx + w^6)(x^3 + x + w^6)$	$y^8 + (w^7x^5 + w^2x^4 + x^3 + wx^2 + w^3x + 2)y^6 + (w^7x^{11} + w^7x^{10} + 2x^9 + wx^8 + x^5 + 2x^4 + w^7x^3 + 2x^2 + 2x + 2)y^4 + (2x^{18} + w^7x^{17} + w^3x^{16} + wx^{15} + w^3x^{14} + w^6x^{13} + w^7x^{12} + w^2x^{11} + w^6x^{10} + 2x^9 + w^5x^7 + w^7x^6 + w^7x^5 + w^2x^2 + 2)y^2 + w^6x^{22} + x^{21} + w^6x^{20} + 2x^{19} + w^6x^{18} + w^5x^{17} + w^7x^{16} + w^6x^{15} + w^3x^{14} + 2x^{13} + wx^{12} + w^6x^{11} + wx^9 + w^6x^8 + w^7x^6 + w^2x^4 + w^5x^3 + 2x^2 + w^6x + 1$	64	$64 - 82$
27	1	$f_1 = x(x + w^{14})$ $f_2 = x^2 + w^2x + w^{10}$	$y^4 + (2x^2 + w^4x + w^{10})y^2 + w^{20}x^2 + w^7x + w^{20}$	38	38
	3	$f_1 = (x + w^5)(x^2 + w^{12}x + w^3)$ $f_2 = x^2 + w^{18}x + w^{25}$	$y^4 + (x^3 + w^3x^2 + wx + w^2)y^2 + x^6 + w^{14}x^5 + 2x^4 + w * x^3 + w^9x^2 + w^8x + 1$	52	56
	4	$f_1 = (x + w)(x^2 + w^2)$ $f_2 = x^3 + w^7x + w^{25}$	$y^4 + (2x^3 + wx^2 + w^{19}x + w^{17})y^2 + w^2x^4 + w^{23}x^3 + 2x^2 + w^2x + w^{12}$	64	$64 - 66$
	5	$f_1 = x(x + w)(x + w^2)(x + w^3)(x + w^{20})(x + w^{22})$ $f_2 = x(x + w)(x + w^2)(x + w^7)(x + w^{22})(x + w^{25})$	$y^4 + (2x^6 + w^6x^5 + w^{23}x^4 + 2x^3 + w^6x^2 + w^{25}x)y^2 + w^4x^{10} + w^9x^9 + w^{22}x^8 + w^{10}x^7 + x^6 + w^{24}x^5 + w^{10}x^4 + w^{20}x^3 + w^{20}x^2$	68	$68 - 76$
	6	$f_1 = x(x + w)(x + w^2)(x + w^4)(x + w^{20})(x + w^{22})$ $f_2 = x(x + w)(x + w^2)(x + w^5)(x + w^{12})(x + w^{24})$	$y^4 + (2x^6 + x^5 + w^4x^4 + w^{18}x^3 + w^{24}x^2 + w^{10}x)y^2 + w^2x^{10} + w^7 * x^9 + w^8x^8 + w^{12}x^7 + w^4x^6 + w^{16}x^5 + w^5x^4 + w^{22}x^3 + w^{24}x^2$	76	$76 - 86$
	7	$f_1 = x(x^3 + w^{19}x^2 + w^{18}x + w^{16})(x^2 + w^2x + w^{22})$ $f_2 = (x^4 + w^7x^2 + x + w^4)(x^2 + w^2x + w^{22})$	$y^4 + (2x^6 + w^7x^5 + w^{22}x^4 + w^6x^3 + wx^2 + w^8x + 1)y^2 + w^{12}x^{10} + w^8x^9 + w^{15}x^8 + x^7 + 2x^6 + w^9x^5 + 2x^3 + w^{17}x^2 + 2x + 1$	**82**	$79 - 96$

$\mathbf{q}$	$\mathbf{g_L}$	$\mathbf{f_i}$	$\mathbf{L = \mathbb{F}_q(x, y)}$	$\mathbf{N(L)}$	
	8	$f_1 = (x + w^{15})(x^2 + w^{25}x + w^5)(x^5 + w^3x^4 + w^{20}x^3 + w^{21}x^2 + w^{11}x + w^{18})$ $f_2 == (x + w^{18})(x^2 + w^2x + w^{22})(x^5 + w^3x^4 + w^{20}x^3 + w^{21}x^2 + w^{11}x + w^{18})$	$y^4 + (2x^8 + wx^7 + w^7x^6 + w^{16}x^5 + w^3x^4 + w^3x^3 + w^{18}x^2 + w^{15} * x + w^{17})y^2 + w^8x^{14} + w^{21}x^{13} + w^{10}x^{12} + w^{23}x^{11} + w^{22}x^{10} + w^{16}x^9 + w^3x^8 + wx^7 + w^{25}x^6 + w^{20}x^5 + w^{15}x^4 + w^{19}x^3 + w^3x^2 + x + w^6$	84	$92 - 106$
	9	$f_1 = x + w^{12}$ $f_2 = x(x^2 + w^8x + w^{18})$ $f_3 = x^3 + w^7x + w^{25}$	$y^8 + (x^3 + w^{21}x^2 + w^7x)y^6 + (x^6 + w^8x^5 + w^5x^4 + w^{25}x^3 + w^6x^2 + w^4x + w^{11})y^4 + (w^{22}x^7 + w^3x^6 + w^{18}x^5 + w^3x^4 + wx^3 + w^8x^2 + w^{15} * x)y^2 + w^2x^8 + w^{14}x^7 + w^{11}x^6 + w^{24}x^4 + w^8x^3 + w^{23}x^2 + w^2x + w^{22}$	88	$91 - 116$
	10	$f_1 = (x + w)(x^2 + w^{12}x + w^{17})(x^5 + w^5x^4 + x^3 + w^{20}x^2 + w^8x + w^2)$ $f_2 = (x + w)(x^2 + w^{12}x + w^{17})(x^5 + w^4x^4 + wx^3 + w^9x^2 + w^9x + 1)$	$y^4 + (2x^8 + w^{23}x^7 + w^{14}x^6 + w^2x^5 + w^8x^4 + w^{12}x^3 + w^5x^2 + w^3x + 2)y^2 + w^{14}x^{14} + w^5x^{13} + w^5x^{12} + w^{21}x^{11} + wx^{10} + w^{25}x^9 + w^{15}x^8 + w^{18}x^7 + w^{24}x^6 + w^{22}x^5 + w^{22}x^4 + w^4x^3 + wx^2 + w^{16}x + w1^8$	**94**	$91 - 126$
	11	$f_1 = (x + w)(x^2 + w^2)$ $f_2 = (x + w^5)(x^2 + w^{12}x + w^3)$ $f_3 = x^2 + w^{18}x + w^{25}$	$y^8 + (x^3 + w^9x^2 + w^{23}x + w^{24})y^6 + (x^6 + w^{14}x^5 + 2x^4 + w^{24}x^3 + w^3x^2 + 2x + 1)y^4 + (x^8 + w^{11}x^7 + w^{25}x^6 + w^{12}x^5 + w^7x^4 + w^6x^3 + w^6x^2 + w^{17}x + w^{17})y^2 + x^{10} + w^8x^9 + w^{21}x^8 + w^{22}x^7 + w^{14}x^6 + w^{23}x^5 + w^7x^4 + w^{21}x^3 + w^7x^2 + w^{18}x + w^6$	**96**	≤ 136
81	1	$f_1 = x^2 + w, f_2 = x^2 + w^{41}$	$y^4 + 2x^2y^2 + w^2$	100	100
	2	$f_1 = (x + w^{57})(x^3 + w^2x + w^6)$ $f_2 = (x + w^{79})(x^3 + w^2x + w^6)$	$y^4 + (2x^4 + w^{24}x^3 + w^{42}x^2 + w^{16}x + w^{30})y^2 + w^2x^6 + w^{44}x^4 + w^{48}x^3 + w^6x^2 + w^{50}x + w^{14}$	116	118
	4		$y^4 + (2x^3 + w^{42}x^2 + w^6x + w^{53})y^2 + w^{48}x^2 + w^{12}x + w^{56}$	150	154
	5	$f_1 = x(x + w^{12})$ $f_2 = (x + w^{50})(x + w^{67})$ $f_3 = (x + w^4)(x + w^{23})$	$y^8 + (w^{50}x + w^7)y^6 + (w^{50}x^3 + w^{22}x^2 + w^{34}x + w^{64})y^4 + (2x^6 + w^{50}x^5 + w^{50}x^4 + w^{27}x^3 + w^8x^2 + w^{26}x + w^{41})y^2 + w^{20}x^6 + w^{55}x^5 + 2x^4 + w^{49}x^3 + w^{25}x^2 + w^4x + w^{68}$	**160**	$156 - 172$

q	$\mathbf{g_L}$	$\mathbf{f_i}$	$\mathbf{L = \mathbb{F}_q(x, y)}$	$\mathbf{N(L)}$	
	7	$f_1 = (x^2 + w)(x^4 + w^4x^2 + w^{13})$ $f_2 = (x^2 + w)(x^4 + w^2x^2 + 2x + w^{22})$	$y^4 + (2x^6 + w^{12}x^4 + 2x^3 + w^{24}x^2 + w^{41}x + w^{26})y^2 + w^{54}x^8 + w^{67}x^7 + w^3x^6 + w^{51}x^5 + w^{73}x^4 + w^{12}x^3 + w^{50}x^2 + w^{68}x + w^{54}$	**172**	$160 - 208$
	13	$f_1 = x^4 + w^7x^2 + w^{77} + w^{16}$ $f_2 = x^4 + w^{15}x^2 + w^{21}x + w^{48}$ $f_3 = x^2 + w^{23} + w^{59}$	$y^8 + (x^4 + w^{39}x^2 + w^{15}x + w^{35})y^6 + (x^8 + w^{27}x^6 + w^9x^5 + w^{57}x^4 + w^{29}x^3 + w^{65}x^2 + w^{10}x + w^{65})y^4 + (x^{10} + w^{23}x^9 + w^{61}x^8 + w^{12}x^7 + w^{11}x^6 + w^6x^5 + w^{22}x^4 + w^{30}x^3 + w^{75}x^2 + w^{38}x + w^{70})y^2 + x^{12} + w^{63}x^{11} + w^{52}x^{10} + w^{46}x^9 + w^{43}x^8 + w^{23}x^7 + w^{68}x^6 + w^{50}x^5 + w^{50}x^4 + w^{75}x^3 + w^{16}x^2 + w^8$	**224**	≤ 315

References

[1] R. Auer. *Ray class fields of global function fields with many rational places.* PhD thesis, Carl-von-Ossietzsky-Universität, Oldenburg, 1999.

[2] I. Duursma, H. Stichtenoth, and C. Voss. Generalized Hamming weights for duals of BCH codes, and maximal algebraic function fields. In *Arithmetic, geometry and coding theory (Luminy, 1993)*, pages 53–65. de Gruyter, Berlin, 1996.

[3] The KANT Group. KANT V4. *J. Symb. Comp.*, 24(3):267–283, 1997.

[4] R. E. MacRae. On unique factorization in certain rings of algebraic functions. *J. Algebra*, 17:243–261, 1971.

[5] M. Rosen. The Hilbert class field in function fields. *Exposition. Math.*, 5(4):365–378, 1987.

[6] H. Stichtenoth. *Algebraic Function Fields and Codes*. Universitext. Springer-Verlag, 1993.

[7] G. van der Geer and M. van der Vlugt. Constructing curves over finite fields with many points by solving linear equations. In *Applications of curves over finite fields (Seattle, WA, 1997)*, pages 41–47. Amer. Math. Soc., Providence, RI, 1999.

[8] G. van der Geer and M. van der Vlugt. New tables for the function $N_q(g)$. *http://www.wins.uva.nl/geer*, 2000.